

Protocol for the Use of the ecSatter Inventory 2.0™

Thank you for your interest in the copyrighted ecSatter Inventory (ecSI 2.0™). Eating competence is theoretically (E. M. Satter, 2007a) and practically (J. S. Krall & Lohse, 2010; E. Satter, 2007; E. M. Satter, 2007b) defined, and is assessed by the ecSI 2.0™. ecSI 2.0™ is valid for use in a mixed population of English-speaking women and men of mixed socio-economic status in the United States age 18 years and above (J.S. Krall, Lohse, & Satter, 2010; Lohse, Satter, Horacek, Gebreselassie, & Oakland, 2007; Lohse, 2015). Testing indicates that high scorers show superior biological health indicators (Lohse et al., 2010; Psota, Lohse, & West, 2007), the same or lower BMIs, (J.S. Krall et al., 2010) better sleep, (Quick et al., 2013) are more active, (Greene et al., 2011) and eat a more healthful diet (Lohse, Bailey, Krall, Wall, & Mitchell, 2012; Lohse et al., 2010). In addition to English, ecSI 2.0™ is available in the following languages: Arabic, Finnish, German, Japanese and Spanish. For each language, the survey has followed the approved translation protocol.

Permission to use the ecSI 2.0™ requires an application process. Usage is available to research projects that contribute toward developing and examining the instrument and furthering the study of the eating competence construct. ecSI 2.0™ tests the degree to which the individual's eating attitudes and behaviors conform to the principles of the Satter Eating Competence Model (ecSatter) (E. M. Satter, 2007a). To gain permission to use ecSI, the research design must test the intervention in a manner that neither conjures false support nor encourages unfounded dismissal of ecSatter.

Possible projects include:

- Surveying eating competence in particular populations and/or validating ecSI 2.0™ with specific populations, such as Spanish-speaking populations, disabled people or people with chronic diseases such as diabetes.
- Large-group surveys that correlate ecSI 2.0™ scores with other parameters such as BMI, as illustrated in the original validation studies, or with metabolic parameters, nutritional status, etc.
- Using ecSI 2.0™ to measure the impact of a *clearly* defined education model or clinical intervention.

Gaining permission *for each intended usage* requires completing an application and will allow us to include you in a community of educators and researchers with expertise in eating competence. Upon application approval, you will receive the usage guidelines.

In summary, here is the process for gaining permission for use ecSI 2.0™.

- 1) Review the [application guidelines](#).
- 2) Submit your application, following the link [here](#) or at the bottom of the application guidelines.
- 3) Upon review of this application, respond to any questions and/or comments. Upon approval, download and use the survey and scoring information as outlined in your proposed project.
- 4) After your project is completed, prepare a confidential outcomes report and upload to [this submission form](#).
- 5) We appreciate you sharing any publications or presentations based on your project with the ecSI 2.0™ community [by uploading them using this submission form](#).

References

- Greene GW, Schembre SM, White AA, Hoerr SL, Lohse B, Shoff S, Horacek T, Riebe D, Patterson J, Phillips BW, Kattelman KK, Blissmer B. Identifying clusters of college students at elevated health risk based on eating and exercise behaviors and psychosocial determinants of body weight. *J Amer Diet Assoc.* 2011;111:394-400.
- Krall JS, Lohse B. Cognitive testing with female nutrition and education assistance program participants informs content validity of the Satter Eating Competence Inventory. *J Nutr Educ Behav.* 2010;42(4):277-283.
- Krall JS, Lohse B, Satter E. Validation of a measure of eating competence for low-income females. International Society for Behavioral Nutrition and Physical Activity Annual Conference Program. 2010; pp. 143-144.
- Lohse B. The Satter eating competence inventory for low-income persons is a valid measure of eating competence for persons of higher socioeconomic position. *Appetite.* 2015;87:223-228.
- Lohse B, Bailey R, Krall J, Wall D, Mitchell DC. Diet quality is related to eating competence in cross-sectional sample of low-income females surveyed in Pennsylvania. *Appetite.* 2012;58(2):645-650.
- Lohse B, Psota T, Estruch R, Zazpe I, Sorli JV, Salas-Salvadó J, Serra M, Krall JS, Márquez F, Ros E on behalf of the PREDIMED study investigators. Eating Competence of Elderly Spanish Adults Is Associated with a Healthy Diet and a Favorable Cardiovascular Disease Risk Profile. *J Nutr.* 2010;140:1322-1327.
- Lohse B, Satter E, Horacek T, Gebreselassie T, Oakland MJ. Measuring eating competence: Psychometric properties and validity of the ecSatter inventory. *J Nutr Educ Behav.* 2007; 39(5S):S154-S166.
- Psota TL, Lohse B, West SG. Associations between eating competence and cardiovascular disease biomarkers. *J Nutr Educ Behav.* 2007;39(5S):S171-S178.
- Quick V, Byrd-Bredbenner C, White AA, Brown O, Colby S, Shoff S, Lohse B, Horacek T, Kidd T, and Greene G.) Eat, sleep, work, play: Associations of weight status and health-related behaviors among young adult college students. *Am J Health Promot.* 2014;29(2):e64-72.
- Satter E. Hierarchy of food needs. *J Nutr Educ Behav.* 2007; 39: S187-188.
- Satter EM. Eating Competence: definition and evidence for the Satter Eating Competence Model. *J Nutr Educ Behav.* 2007a; 39: S142-S153.
- Satter EM. Nutrition education with the Satter Eating Competence Model. *J Nutr Educ Behav.* 2007b; 39: S189-S194.